



Algumas propriedades das funções trigonométricas

1. $\forall a \in \mathbb{R} \quad \sin^2 a + \cos^2 a = 1$
2. $\forall a \in \mathbb{R} \setminus \{\frac{\pi}{2} + k\pi : k \in \mathbb{Z}\} \quad 1 + \tan^2 a = \sec^2 a$
3. $\forall a \in \mathbb{R} \setminus \{k\pi : k \in \mathbb{Z}\} \quad 1 + \cot^2 a = \operatorname{cosec}^2 a$
4. $\forall a \in \mathbb{R} \quad \sin(-a) = -\sin a \quad (\sin \text{ é ímpar})$
5. $\forall a \in \mathbb{R} \quad \cos(-a) = \cos a \quad (\cos \text{ é par})$
6. $\forall a \in \mathbb{R} \quad \cos(\frac{\pi}{2} - a) = \sin a \quad \text{e} \quad \sin(\frac{\pi}{2} - a) = \cos a$
7. $\forall a \in \mathbb{R} \quad \sin(a + 2\pi) = \sin a \quad (\sin \text{ tem período } 2\pi)$
8. $\forall a \in \mathbb{R} \quad \cos(a + 2\pi) = \cos a \quad (\cos \text{ tem período } 2\pi)$
9. $\forall a, b \in \mathbb{R} \quad \sin(a + b) = \sin a \cos b + \sin b \cos a$
10. $\forall a, b \in \mathbb{R} \quad \cos(a + b) = \cos a \cos b - \sin a \sin b$
11. $\forall a, b \in \mathbb{R} \quad \cos a - \cos b = -2 \sin \frac{a-b}{2} \sin \frac{a+b}{2}$
12. $\forall a, b \in \mathbb{R} \quad \sin a - \sin b = 2 \cos \frac{a-b}{2} \sin \frac{a+b}{2}$

Algumas propriedades das funções hiperbólicas

$$\begin{array}{rcl} \operatorname{sh} : & \mathbb{R} & \longrightarrow & \mathbb{R} \\ & x & \longmapsto & \frac{e^x - e^{-x}}{2} \end{array}$$

$$\begin{array}{rcl} \operatorname{ch} : & \mathbb{R} & \longrightarrow & \mathbb{R} \\ & x & \longmapsto & \frac{e^x + e^{-x}}{2} \end{array}$$

1. $\forall a \in \mathbb{R} \quad \operatorname{ch}^2 a - \operatorname{sh}^2 a = 1$
2. $\forall a \in \mathbb{R} \quad \operatorname{th}^2 a + \operatorname{sech}^2 a = 1$
3. $\forall a \in \mathbb{R} \setminus \{0\} \quad \operatorname{coth}^2 a - \operatorname{cosech}^2 a = 1$
4. $\forall a \in \mathbb{R} \quad \operatorname{sh}(-a) = -\operatorname{sh} a \quad (\operatorname{sh} \text{ é ímpar})$
5. $\forall a \in \mathbb{R} \quad \operatorname{ch}(-a) = \operatorname{ch} a \quad (\operatorname{ch} \text{ é par})$
6. $\forall a, b \in \mathbb{R} \quad \operatorname{sh}(a + b) = \operatorname{sh} a \operatorname{ch} b + \operatorname{sh} b \operatorname{ch} a$
7. $\forall a, b \in \mathbb{R} \quad \operatorname{ch}(a + b) = \operatorname{ch} a \operatorname{ch} b + \operatorname{sh} b \operatorname{sh} a$
8. $\forall n \in \mathbb{N} \quad \forall a \in \mathbb{R} \quad (\operatorname{ch} a + \operatorname{sh} a)^n = \operatorname{ch}(na) + \operatorname{sh}(na)$